

Lactobacillus reuteri Tablets Suppress *Helicobacter pylori* Infection
—A Double-blind Randomised Placebo-controlled
Cross-over Clinical Study—

Kyoto IMASE, Akifumi TANAKA, Kengo TOKUNAGA, Hajime SUGANO, Hitoshi ISHIDA
& Shin'ichi TAKAHASHI

Third Department of Internal Medicine, Kyorin University School of Medicine

(Received : October 16, 2006)

(Accepted : March 20, 2007)

Key words : *Helicobacter pylori*, probiotics, *Lactobacillus reuteri*, randomised controlled study

Abstract

We studied the effect of *Lactobacillus reuteri* strain SD2112 Tablets Reuterina (ERINA Co., Inc.), in suppressing *H. pylori* urease activity and to use the urea breath test (UBT) as a marker for the burden of infection. Method 1 : Assessment of UBT and *H. pylori* density. Subjects were 33 *H. pylori*-positive patients from whom were obtained gastric biopsy specimens by upper gastrointestinal endoscopy. The correlation between UBT and *H. pylori* density was investigated. Individual UBT was established for each patient. Patients were divided by *H. pylori* density was 3 groups : Group I (low-density), Group II (moderate-density), and Group III (high-density). The individual UBTs were then correlated to the established *H. pylori* quantity. Method 2 : Assessment of suppressive effect of *L. reuteri* on *H. pylori* urease activity. Subjects were 40 asymptomatic volunteers with an UBT exceeding 15%, randomly allocated to four groups : Subjects in Group A underwent active treatment for 4 weeks (period 1) and placebo treatment for the following 4 weeks (period 2). These in Group B underwent treatment in reverse order. Those in Group C underwent placebo. Group D consisted of volunteers with negative UBT undergoing active treatment for the full 8 weeks. Result 1 : UBT was $11.6 \pm 2.0\%$, $22.1 \pm 2.6\%$, and $35.4 \pm 7.6\%$ in Groups I, II, and III, showing UBT that increased significantly (I vs. II : $p < 0.01$ and I vs. III : $p < 0.05$) based on *H. pylori* density. Result 2 : Significant differences were seen in the decrease in UBT before versus after medication in Groups A and B. In Group A, lower UBT was maintained until the end of the full 8-week period. The overall decrease in UBT due to medication with *L. reuteri* Tablets was $69.7 \pm 4.0\%$ ($p < 0.05$). Conclusion : Administration of *L. reuteri* Tablets [Reuterina (ERINA Co., Inc.)] significantly decreased UBT in *H. pylori*-positive subjects, demonstrating that *L. reuteri* suppresses *H. pylori* urease activity and *H. pylori* density.

[J.J.A. Inf. D. 81 : 387~393, 2007]

Introduction

Helicobacter pylori has been recognized as a cause of gastric mucosal injury. The treatment of first choice in the management of *H. pylori*-positive peptic ulcer is eradication of the organism, but this fails, in 10-20% of patients undergoing eradication^{1)~5)}. A decline in eradication due to increased drug-resistant *H. pylori* has become a

Correspondence to : Kyoto IMASE

Third Department of Internal Medicine, Kyorin University School of Medicine, 16-20-2 Shinkawa, Mitaka-shi, Tokyo 181-8611, Japan

平成19年7月20日

worldwide issue. No definitive treatment policy for refractory cases, in whom eradication of *H. pylori* has failed, has been established. A large number of asymptomatic *H. pylori*-infected subjects have no clinical need for eradication, yet, this group of "healthy" subjects should be considered at risk to develop disease if infection remains untreated⁶⁾. Probiotic treatments designed to improve disordered indigenous intestinal bacterial flora are drawing increased attention. Of these, lactic acid bacteria have been reported to suppress *H. pylori* gastritis by reducing *H. pylori* density. We studied the effect of one of these probiotics, *Lactobacillus reuteri* strain SD2112 Tablets [*L. reuteri* tablet, Reuterina (ERINA Co., Inc.)], in suppressing *H. pylori* urease activity and used the urea breath test (UBT) as a marker for the burden of infection.

Materials and Methods

1. UBT assessment

UBT was used for diagnosing *H. pylori* infection. Fasting subjects were instructed to orally take a ¹³C-urea preparation (UBIT Tablet 100mg, Otsuka Pharmaceutical Co., Ltd., Tokyo, Japan), in the early morning with 100 mL of water while sitting. Exhalation specimens were collected on two occasions, before and 20 minutes after oral administration and UBT was conducted with ¹³C-urea measurement (UbiT-IR300; Otsuka Electronics Co., Ltd., Osaka, Japan). The formula used to calculate ¹³C-urea was: $\Delta^{13}\text{C} = \delta^{13}\text{C}$ (after administration of UBIT Tablets) - $\delta^{13}\text{C}$ (before administration of UBIT Tablets). Based on the results of large-scale prospective clinical trials in Japan⁷⁾, a 2.5% cut off value for UBT was used. UBT exceeding 2.5% was taken as positive. It was shown elsewhere⁸⁾ that the reproducibility of UBT increases with time. Detailed assessment of this report revealed a trend for UBT to be unstable at or above 50%. Another report⁹⁾ has shown marked variation in urease activity not be attributable to technical diagnostic reasons, although instability tended to occur when urease activity was high. Based on these reports, we studied UBT 8 weeks before medication and established UBT value as a standard, compared to UBT, the standard vs before medication. Subjects in whom the difference in UBT standard versus before was 50% or higher were excluded from the present analysis because of low reliability.


2. Correlation of UBT to *H. pylori* density

Informed consent was obtained from 33 *H. pylori*-positive patients (age: 28-63 years; mean: 49.2±9.90 years; 30 males and 3 females) undergoing upper gastrointestinal endoscopy at Kyorin University Hospital. Gastric biopsy specimens were obtained and individual UBT was established for each patient. Biopsy samples were collected from two sites on the greater gastric curvature, the antrum and the upper corpus. Each specimen was immediately embedded in Seed Tube HP (Eiken Chemical, Tokyo, Japan) and stored at 4°C. Each biopsy specimen was placed in 1mL of buffered sodium chloride peptone solution (Nissui, Tokyo, Japan), and inoculated onto Skirrow's media (Kyokuto Pharmaceutical Industrial Co. Ltd., Tokyo, Japan) at 37°C for 5-7 days under microaerobic conditions (5% O₂, 10% CO₂, 85% N₂). The amount of *H. pylori* was expressed as described elsewhere¹⁰⁾. The correlation between UBT and bacterial density was investigated. Patients were divided into three groups by *H. pylori* density: Group I [less than 1×10⁵ colony forming units (cfu) /mL], Group II (between 1×10⁵ and up to 5×10⁵ cfu/mL), and Group III (more than 5×10⁵ cfu/mL). Patients in these groups were matched for age, gender and disease. The individual UBT was then correlated to the established *H. pylori* density.

3. Assessment of suppressive effect of *L. reuteri* on *H. pylori* urease activity

Informed consent was obtained from 179 asymptomatic volunteers (age: 18-65 years; mean: 44.5±10.5 years; 84 males and 95 females). Of 40 volunteers in the study, 5 were *H. pylori*-negative and 35 were *H. pylori*-positive (47.8±10.7 years; 19 males and 21 females). We studied UBT four times; 8 weeks before, before, 4 weeks after and 8 weeks after medication. Volunteers with an UBT exceeding 15% were randomly allocated to either Group A, B or C. Group A (n=15, 48.1±13.5 years) and Group B (n=15, 48.9±7.05 years) entered a randomised, double-blind crossover study where the effects of *L. reuteri* tablets (active ingredient 10⁸ Colony Forming Units of *L. reuteri*, strain ATCC 55730) was compared to the effect of identical placebo tablets. Placebo tablets contained no *L. reuteri*, and the isomalt content was slightly higher to compensate for weight. The daily dose was 4 tablets. Group A received active treatment for 4 weeks (period 1) and placebo treatment for the following 4 weeks (Period 2). Subjects in Group B received the study treatment in the reverse order, i.e., placebo tablets for 4 weeks

Fig. 1 Correlation between UBT and *H.pylori* density
* p < 0.05 (Wilcoxon signed-ranks' test)



(Period 1) and *L. reuteri* tablets during the following 4 weeks (Period 2). Group C (n=5, 48.8±11.6 years) was also part of the randomised double-blind study, but received placebo for the full 8 weeks. The purpose of this group was to assess the natural intraindividual variation in UBT measurement over time. To measure true variations in UBT, volunteers had to have an initial UBT of 15% or more. Group D (n=5, 42.6±9.39 years) consisted of volunteers with a negative UBT and were treated for the full 8 weeks with 4 tablets daily of *L. reuteri* tablets. The purpose of this group was to ascertain that *L. reuteri* per se did not influence UBT assessment. Exclusion criteria were chronic diseases, those receiving oral medications such as antac-

ids and antibiotics; yogurt consumption, and participation in other clinical studies within the preceding 4-week period.

4. Analysis

All data obtained was expressed as mean standard error of the mean (SEM). The significance of difference between groups was analyzed using the Wilcoxon test, and software used was Stat view ver. 5.0 for Macintosh. The study protocol was submitted to the Kyorin University Ethical Committee for review and approval.

Results

1. UBT and *H. pylori* density

As shown in Figure 1, UBT was 11.6±2.0% (mean±SEM), 22.1±2.6% and 35.4±7.6% in Groups I (low density group: $0.4 \times 10^5 \pm 0.1 \times 10^5$ cfu/mL), II (moderate density group: $2.5 \times 10^5 \pm 0.4 \times 10^5$ cfu/mL) and III (high density group: $8.8 \times 10^5 \pm 0.7 \times 10^5$ cfu/mL), showing that UBT increased significantly (I vs. II: p<0.01 and I vs. III: p<0.05) with to bacterial density.

2. Suppressive effect of *L. reuteri* Tablets on *H. pylori*

(a) Variation in UBT

As shown in Figures 2a through -2d, significant (p<0.05) differences were seen in the decrease in UBT before versus after 4 weeks of medication in Group A (n=13) and before versus after 8 weeks of medication in Group B (n=11), whereas no significant difference was seen in Group C (n=5) or D (n=5).

(b) UBT decrease

The decrease in UBT during the period that subjects took *L. reuteri* Tablets was studied in 24 subjects in Groups A and B [6 subjects were disqualified and excluded from the study]. Decrease was assessed based on UBT before and after 4 weeks of medication in Group A and before and after 8 weeks of medication in Group B. In the comparison with before medication, UBT changed 65.3±5.8% after 4 weeks of medication (p<0.05) in Group A, and 74.9±5.2% after 8 weeks of medication (p<0.05) in Group B, i.e., UBT differed significantly. No significant difference was seen in the decrease in Group C or D. Figure 3 shows UBT variation during oral medication with *L. reuteri* Tablets in all Group A and B subjects. The overall decrease in UBT due to medication with *L. reuteri* Tablets was 69.7±4.0% (p<0.05) during this period, i.e., the difference was significant.

(c) Number of individuals responding to treatment.

All in all, 19 of the 24 subjects responded favourably to treatment with *L. reuteri* tablets whereas only 10 responded to placebo treatment (p<0.01, Fisher's exact test). It is noteworthy that 7 of the 10 who responded favourably to the placebo treatment belonged Group A, indicative that they could be "late" responders to *L. reuteri*.

Discussion

There are arguments for and against the correlation between *H. pylori* density and UBT. Some reports¹¹⁾¹²⁾

Fig. 2 Changes in UBT in Group A (a), B (b), C (c) and D (d). UBT treatment differs in Group A and B. For Group A (n = 13), UBT was before versus 4 weeks after medication. For Group B (n = 11), UBT was before versus 8 weeks after medication. Group C (n = 5), and Group D (n = 5).





Fig. 3 Reduction in UBT from the beginning of oral medication (n = 24) with *L. reuteri* Tablets to the end in all subjects.



of *H. pylori* and the severity of gastritis. We showed a significant correlation between *H. pylori* density and the gastritis score¹⁰. When *H. pylori* density increased, the pathogen also presumably increased and induced gastric mucosal injury in the host.

Probiotics are increasingly being used as dietary supplements, to promote health, and to treat of digestive organ diseases^{16,17}. *Lactobacillus gasseri* OLL 2716 (LG21), a probiotic, is reported to decrease UBT, the pepsinogen I to II ratio, and serum *H. pylori* antibody titer in *H. pylori*-positive patients¹⁸. LG21 is believed to suppress *H. py-*

have indicated a significant correlation between *H. pylori* density determined histologically and UBT, while others⁹ have shown no correlation between these factors. We compared the possible correlation between *H. pylori* density and UBT in three groups with different bacterial densities, i.e., low, moderate, and high. As shown in Figure 1, a significant difference was seen in UBT among these three groups, suggesting a correlation between bacterial density and UBT. Regarding the etiology of gastric mucosal injury, some reports^{13,14} have indicated that ammonia is produced by *H. pylori* urease activity and that reversed hydrogen ion diffusion produces associated injury. Another report⁹ discussing the correlation between *H. pylori* urease activity and the gastritis score, while yet another¹⁵ showed that UBT is useful for evaluating density and activity

lori density. We used UBT to assess the suppressive effect of *L. reuteri* on *H. pylori* in healthy volunteers. As shown in Figure 2a, administering *L. reuteri* Tablets in Group A resulted in significantly decreased UBT at the end of the first-4 week period and this lower UBT value was maintained until the end of the full 8-week period, during the latter half of which subjects took placebo Tablets. In Group B, UBT decreased during *L. reuteri* administration (Figure 2b). Consistent with these results, some reports¹⁹⁾²⁰⁾ have shown that *L. johnsonii* La1 strain (*Lactobacillus*), administered to mice, survived for a long time in the intestine even after administration was discontinued. In the present study, *L. reuteri* showed prolonged survival in the stomach and the intestine²¹⁾ even after the administration of *L. reuteri* Tablets was discontinued.

L. reuteri, which is a common intestinal bacterium in vertebrates, is known to be nonpathogenic and very safe. It is reported to produce an antibiotic-like substance and to adhere to the digestive tract, suppressing the attachment of pathogenic bacteria to the digestive tract mucosa²²⁾. This antibiotic-like substance, reuterin (3-hydroxypropionate aldehyde), is from anaerobically growing organisms and is produced in the presence of glycerol²³⁾. It has a broad antibacterial spectrum, which includes Gram-positive bacteria, Gram-negative bacteria, yeasts and other fungi, according to one report²⁴⁾. It is reported²⁵⁾ that *L. reuteri* inhibits *in vitro* attachment of *H. pylori* to the gastric mucosa and that *L. reuteri* inhibits the growth of *H. pylori in vitro*. Decreased UBT with *L. reuteri* administration, in the present study, may be explained by an action of reuterin on *H. pylori* involving indirect inhibition of attachment of *H. pylori* to the gastric mucosa, suppressing of *H. pylori* density.

H. pylori infection is known to stimulate both the gastric mucosa and lymphocytes, promoting the secretion of cytokines including interleukin (IL)-8 and TNF- α , and to cause gastritis. *H. pylori* infection induces inflammation in gastric mucosa and that causes gastritis. In the above discussion on LG21, IL-8 production was suppressed by LG21 *in vitro*. There are also reports²⁶⁾ showing that *Lactobacillus* suppresses these cytokines and ameliorates inflammatory intestinal diseases and allergic reactions. *L. reuteri* ATCC 55730 given exogenously in tablet form leads to a significant colonisation of the human gastric antrum²¹⁾, the major site of *H. pylori* infection. Thus, a close interaction between *L. reuteri* and the pathogen can be expected *in vivo* after administration of the probiotic. These reports raise the possibility that *L. reuteri* Tablets suppress *H. pylori*-derived gastritis by suppressing gastric mucosal cytokines.

Conclusions

Administration of *L. reuteri* Tablets [Reuterina (ERINA Co.,Inc.)] significantly decreased UBT in *H. pylori*-positive subjects, demonstrating that *L. reuteri* suppresses *H. pylori* density. It is hoped that, *L. reuteri* Tablets can be used both to prevent asymptomatic *H. pylori* positive subjects to develop clinical symptoms and to improve gastritis, and also improve symptom and gastritis in *H. pylori* infected patients, in whom *H. pylori* eradication has failed.

Acknowledgement

We thank ERINA Co.,Inc. for financial support for this study and Anders Zachrisson (BioGaia AB) advice. We also thank study, participants without who this study could not have been done.

References

- 1) Miwa H, Ohkura R, Murai T, Sato K, Nagahara A, Hirai S, *et al.* : Impact of rabeprazole, a new proton pump inhibitor, in triple therapy for *Helicobacter pylori* infection-comparison with omeprazole and lansoprazole. *Aliment Pharmacol Ther* 1999 ; 13 : 741—6.
- 2) Asaka M, Sugiyama T, Kato M, Satoh K, Kuwayama H, Fukuda Y, *et al.* : A multicenter, double-blind study on triple therapy with lansoprazole, amoxicillin and clarithromycin for eradication of *Helicobacter pylori* in Japanese peptic ulcer patients. *Helicobacter* 2001 ; 6 : 254—61.
- 3) Miwa H, Nagahara A, Sato K, Ohkura R, Murai T, Shimizu H, *et al.* : Efficacy of 1 week omeprazole or lansoprazole-amoxicillin-clarithromycin therapy for *Helicobacter pylori* infection in the Japanese population. *J Gastroenterol Hepatol* 1999 ; 14 : 317—21.
- 4) Miwa H, Ohkura R, Murai T, Nagahara A, Yamada T, Ogihara T, *et al.* : Effectiveness of omeprazole-amoxicillin-

- clarithromycin (OAC) therapy for *Helicobacter pylori* infection in a Japanese population. *Helicobacter* 1998 ; 3 : 132—8.
- 5) Suzuki J, Mine T, Kobayashi I, Fujita T : Assessment of a new triple agent regimen for the eradication of *Helicobacter pylori* and the nature of *H. pylori* resistance to this therapy in Japan. *Helicobacter* 1998 ; 3 : 59—63.
 - 6) Uemura N, Okamoto S, Yamamoto S : Helocobacter pylori. Infection and the development of gastric cancer. *N Engl J Med* 2001 ; 345 : 829—32.
 - 7) Ohara S, Kato M, Asaka M, Toyota T : Studies of ¹³C-urea breath test for diagnosis of *Helicobacter pylori* infection in Japan. *J Gastroenterol* 1998 ; 33 : 6—13.
 - 8) Logan RP, Polson RJ, Misiewicz JJ, Rao G, Karim NQ, Newell D, *et al.* : Simplified single sample ¹³Carbon urea breath test for *Helicobacter pylori* : comparison with histology, culture, and ELISA serology. *Gut* 1991 ; 32 : 1461—4.
 - 9) Graham DY, Klein PD, Evans DJ Jr, Evans DG, Alpert LC, Opekun AR, *et al.* : *Campylobacter pylori* detected noninvasively by the ¹³C-urea breath test. *Lancet* 1987 ; 23 : 1174—7.
 - 10) Takahashi S, Igarashi H, Ishiyama N, Nakano M, Ozaki M, Ito M, *et al.* : Serial change of gastric mucosa after challenging with *Helicobacter pylori* in the cynomolgus monkey. *Zentralbl Bakteriol* 1993 ; 280 : 51—7.
 - 11) Kobayashi D, Eishi Y, Ohkusa T, Ishige I, Suzuki T, Minami J, *et al.* : Gastric mucosal density of *Helicobacter pylori* estimated by real-time PCR compared with results of urea breath test and histological grading. *J Med Microbiol* 2002 ; 51 : 305—11.
 - 12) Chen X, Haruma K, Kamada T, Mihara M, Komoto K, Yoshihara M, *et al.* : Factors that affect results of the ¹³C urea breath test in Japanese patients. *Helicobacter* 2000 ; 5 : 98—103.
 - 13) Hazell SL, Lee A : *Campylobacter pyloridis*, urease, hydrogen ion back diffusion, and gastric ulcers. *Lancet* 1986 ; 5 : 15—7.
 - 14) Thomsen L, Tasman-Jones C, Morris A, Wiggins P, Lee S, Forlong C : Ammonia produced by *Campylobacter pylori* neutralizes H⁺ moving through gastric mucus. *Scand J Gastroenterol* 1989 ; 24 : 761—8.
 - 15) Chang YW, Min SK, Kim KJ, Han YS, Lee JH, Dong SH, *et al.* : Delta (¹³C)-urea breath test value is a useful indicator for *Helicobacter pylori* eradication in patients with functional dyspepsia. *J Gastroenterol Hepatol* 2003 ; 18 : 726—31.
 - 16) McGroarty JA : Probiotic use of lactobacilli in the human female urogenital tract. *FEMS Immunol Med Microbiol* 1993 ; 6 : 251—64.
 - 17) Reid G, Bruce AW, McGroarty JA, Cheng KJ, Costerton JW : Is there a role for lactobacilli in prevention of urogenital and intestinal infections? *Clin Microbiol Rev* 1990 ; 3 : 335—44.
 - 18) Sakamoto I, Igarashi M, Kimura K, Takagi A, Miwa T, Koga Y : Suppressive effect of *Lactobacillus gasseri* OLL 2716 (LG21) on *Helicobacter pylori* infection in humans. *J Antimicrob Chemother* 2001 ; 47 : 709—10.
 - 19) Michetti P, Dorta G, Wiesel PH, Brassart D, Verdu E, Herranz M, *et al.* : Effect of whey-based culture supernatant of *Lactobacillus acidophilus* (*johnsonii*) La1 on *Helicobacter pylori* infection in humans. *Digestion* 1999 ; 60 : 203—9.
 - 20) Felley CP, Corthesy-Theulaz I, Rivero JL, Sipponen P, Kaufmann M, Bauerfeind P, *et al.* : Favourable effect of an acidified milk (LC-1) on *Helicobacter pylori* gastritis in man. *Eur J Gastroenterol Hepatol* 2001 ; 13 : 25—9.
 - 21) Valeur N, Engel P, Carbajal N, Connolly E, Ladefoged K : Colonization and immunomodulation by *Lactobacillus reuteri* ATCC 55730 in the human gastrointestinal tract *Appl Environ Microbiol* 2004 ; 70 : 1176—81.
 - 22) Mukai T, Asasaka T, Sato E, Mori K, Matsumoto M, Ohori H : Inhibition of binding of *Helicobacter pylori* to the glycolipid receptors by probiotic *Lactobacillus reuteri*. *FEMS Immunol Med Microbiol* 2002 ; 32 : 105—10.
 - 23) Talarico TL, Casas IA, Chung TC, Dobrogosz WJ : Production and isolation of reuterin, a growth inhibitor produced by *Lactobacillus reuteri*. *Antimicrob Agents Chemother* 1988 ; 32 : 1854—8.
 - 24) Talarico TL, Dobrogosz WJ : Chemical characterization of an antimicrobial substance produced by *Lactobacillus reuteri*. *Antimicrob Agents Chemother* 1989 ; 33 : 674—9.
 - 25) Johnson C, Dicksved J, Jonsson H, Roos S : Anti *Helicobacter pylori* activity among lactic acid bacteria isolated from gastric biopsies and strains of *Lactobacillus reuteri*. *Helicobacter* 2003 ; 8 : 473.
 - 26) Wu J, Wang XX, Takasaki M, Ohta A, Higuchi M, Ishimi Y : Cooperative effects of exercise training and genistein administration on bone mass in ovariectomized mice. *J Bone Miner Res* 2001 ; 16 : 1829—36.

Lactobacillus reuteri 含有プロバイオティクスの *Helicobacter pylori* 抑制効果に関する検討

杏林大学医学部第3内科

今瀬 教人 田中 昭文 徳永 健吾
菅野 朝 石田 均 高橋 信一

この研究は、*Lactobacillus reuteri* strain SD 2112 株 (*L. reuteri*) の *H. pylori* 抑制効果の検討である。今回、我々は *L. reuteri* 含有整腸剤 [*L. reuteri* 錠, Reuterina (ERINA CO., INC.)] の効果について検討を行った。【検討1】 *H. pylori* 菌量と UBT 値の相関：*H. pylori* 陽性消化性潰瘍患者の 33 名を、Group I (低菌量群)、Group II (中菌量群)、Group III (高菌量群) の 3 群に分類し、*H. pylori* 培養と同時に尿素呼気試験 (UBT) を行った。【検討2】 *L. reuteri* の *H. pylori* 抑制効果：*H. pylori* 陽性者 35 名と陰性者 5 名に、二重盲検法による交差試験で *L. reuteri* 錠または placebo 錠を内服させた。UBT は基準値として測定した内服開始 8 週間前の値の他、内服開始前、開始後 4 週間および 8 週間に検査した。【成績1】 *H. pylori* 菌量と UBT 値の間には、正の相関が見られ、Group I と Group III で有意差を認めた ($p < 0.05$)。【成績2】 UBT は、*L. reuteri* 錠の内服期間 (A 群 + B 群) で、前値 100% から $69.7 \pm 4.0\%$ ($p < 0.05$) と有意に減少した。一方で、C 群 (placebo 内服群)・D 群 (*H. pylori* 陰性 *L. reuteri* 錠内服群) では変動は認めなかった。【結論】 *L. reuteri* 含有整腸剤の内服により、UBT 値は有意に減少した。また、*L. reuteri* は *H. pylori* 菌量も減少させる可能性が示唆された。