

第 94 卷 索 引

I		community-based hospital	553
		compliance	654
16SrDNA sequence	325	coronavirus	850
A		COVID-19	483, 490, 495, 500, 507, 514, 520, 528, 535, 542, 545, 549, 553, 563, 568, 573, 579, 583, 587, 591, 596, 600, 604, 608, 814, 821, 855
abscess	662	critical pneumonia	528
acute pulmonary embolism	604	cruise ship	535
acute respiratory disorder	563	D	
acute respiratory distress syndrome	114, 553	d-dimer	604
alkaptonuria	834	discharge	579
allogeneic hematopoietic stem cell transplanta- tion	109	E	
amoebic liver abscess	102	elderly	568, 573
AmpC	310	emergency room (ER)	135
antibiotic prophylaxis	654	enrichment broth	654
antibody	495, 821	<i>Entamoeba histolytica</i>	102
antimicrobial prescription	304	Enterobacteriaceae	310
antimicrobial resistance (AMR) action plan	304	environmental improvement	135
antimicrobial stewardship	304	epidemic	181
antimicrobial susceptibility	186, 310	extended-spectrum β -lactamase (ESBL)	310
asymptomatic carrier	535, 583	extracorporeal membrane oxygenation (ECMO)	528
azithromycin	600	F	
B		false positive	97
bacteremia	844	false-negative	542
birth cohort	174	fatal	672
blood culture	109	febrile neutropenia	109
brain abscess	332	fluoroquinolone	109
C		food	186
catheter-related blood stream infection	321	G	
cerebrospinal fluid	193	genetic testing	297
chest CT	507, 520, 535, 549, 814	ground-glass opacity	507
chest X-ray	520	group B <i>Streptococcus</i>	654
chronologic in CT findings	814		
<i>Citrobacter koseri</i>	834		
clinical course	500, 514, 520		
community acquired pneumonia	507		

H			
HBs antigen	97	mother-to-child transmission	654
health care facility	573	MRSA	186
hemagglutinin-neuraminidase sequence	86	multilocus sequence typing (MLST)	186
hemodialysis	596	multiplex PCR	317
Hickman catheter	321	multiplex RT-nested PCR	86
high sensitivity	97	<i>Mycobacterium haemophilum</i>	325
high-risk exposure	855	<i>Mycobacterium leprae</i>	325
household infection	573	N	
human astrovirus	808	nasal swab	542, 545
human parainfluenza virus	86	nasopharyngeal swab	587
human-to-human transmission		natural course	528, 568
	855	necrotizing soft tissue infection	
hydroxychloroquine	596, 600		834
hyperviscous <i>Klebsiella</i>	839	neonatal fever	317
I		nested multiplex PCR	850
immunochromatography	495	no treatment	568
indirect haemagglutination test		nosocomial infection	553
	102	O	
infant	181	occupational infection	135
infection control	821	odontogenic infection	114
infection prevention	135	outbreak	808
infective endocarditis	667	P	
influenza vaccine	647	PCR	193, 338, 500,
intrapartum	654		545, 579, 583, 587, 591
K		PCR negative	514
Kanagawa model	490	pediatric clinic	304
L		penicillin-resistance	667
leptospirosis	193	phylogenetic analysis	86
levofloxacin	828	positive	579
liver cirrhosis	672	prosthetic joint infection	338
low risk patient	549	<i>Pseudomonas aeruginosa</i>	338
lower respiratory specimen	542	pyomyositis	662
lumbago	608	Q	
lung abscess	839	questionnaire	297
M		R	
mediastinal emphysema	604	rapid diagnosis	317, 495
medical disintegration	490	repeat vaccination	647
melting temperature mapping method	332	resistance	828
mild to moderate case	500	respiratory syncytial virus	181
		<i>Rhizobium radiobacter</i>	321

risk factor	483, 563	<i>Streptococcus salivarius</i>	667
reversetranscriptase-PCR (RT-PCR)	514	string test	839
rubella	174	symptom	591
S		T	
SARS-CoV-2	608, 821	test-negative design	647
seroepidemiology	174	tuberculosis	828
seroprevalence	821	typical symptom	549
serotype 8	808	U	
severe fever with thrombocytopenia syndrome	672	urinary tract infection	310
severe pneumonia	483	V	
severity classification	490	vaccination	135, 174
sinusitis	583	ventricular fibrillation	600
smoking history	483	<i>Vibrio cholerae</i> non-O1 non-O139	
sputum	545, 591	viral load	587
staphylococcal cassette chromosome <i>mec</i> (SCC <i>mec</i>)	186	viral pneumonia	563, 850
<i>Staphylococcus schleiferi</i>	662	virulence factor	844
<i>Streptococcus intermedius</i>	332		
